DANGER

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Foli-Zyme®

PHYSIOLOGICAL SUPPORT

(15 – 0 – 3 – 0)

A liquid fertiliser formulated to enhance healthy growth and assist plants in stress recovery.

ANALYSIS (% W/V):	Nitrogen (as total)	15.0 %
	Potassium (as total)	3.0 %
	Calcium (as total)	3.9 %
	Magnesium (as total)	1.2 %
	Boron (B)	0.25 %
	Cobalt (Co)	0.01 %
	Copper (Cu)	0.14 %
	Manganese (Mn)	0.1 %
	Molybdenum (Mo)	0.024 %
	Zinc (Zn)	0.9 %
	Maximum biuret	1.25 g/kg N

Foli-Zyme® provides a balanced source of nutrition that help a plant overcome the negative impact of stress events.

SHAKE WELL BEFORE USING

Pack sizes: 20, 1000 L

Manufactured by:
Stoller Australia Pty Ltd
1 Creswell Road
Largs Bay, SA, 5016
Australia

Distributed by:
Corteva Agriscience Australia Pty Ltd
Level 9, 67 Albert Avenue
Chatswood, NSW, 2067
Australia

EMERGENCY RESPONSE (ALL HOURS)
RING FROM ANYWHERE IN AUSTRALIA
1800 370 754
(LOCAL CALL FEE ONLY)

IN A TRANSPORT EMERGENCY ONLY
DIAL 000
FOR POLICE OR FIRE BRIGADE

This label complies with the National Code of Practice for Fertiliser Description and Labelling.

Lot Number: See packaging
Manufacturing date: See packaging
Expiration Date: See packaging

™® Trademarks of Corteva Agriscience and its affiliated companies. © 2026 Corteva

™® Trademarks of Corteva Agriscience and its affiliated companies. © 2026 Corteva

DIRECTIONS FOR USE

RESTRAINTS

Excessive use of molybdenum (Mo) can be harmful to stock. Plant levels of molybdenum can be high for 4 weeks after application. It is advisable to keep stock off treated areas for this period. Molybdenum application may include copper deficiency in grazing animals. This is most likely to occur on sandy soils low in copper. Check rate and frequency of molybdenum use with appropriate authorities.

Crop	Rate	Comment
Trees, Fruit, Vines & Nuts	5 - 9 L/ha	Appy 5 - 6 times annually, beginning with delayed dormant spray. Spray intervals of 14 days are suggested for best results. Avoid periods when fruit is extremely sensitive to spray damage.
Citrus	5 - 9 L/ha	Apply every 30 days. Do not mix with petroleum spray oils or ammoniated copper compounds
Grain and cotton crops	5 - 9 L/ha	Apply when crop is 9 cm high. Continue on 14-day intervals until crop is 14-days from harvest.
Field row crops and Vegetables	5 - 10 L/ha	To supplement soil applied fertiliser, use a row directed spray. Commence application when the crop has 2- 3 leaves or is 20 days old. Continue sprays at 10 - 14 day intervals.
Potatoes	10 L/ha	
Lucerne	2.5 - 5 L/ha	Apply 7 – 14 days after cutting to enhance growth and dry matter.
Greenhouse	250 mL/100 L	Apply every 10 to 14 days at first signs of stress.
Turf	500 mL/1000 square metres	Repeat every 2 – 3 weeks until dormancy.

For further information, consult your Corteva Agriscience representative for advice specific to your situation.

GENERAL INSTRUCTIONS

Foli-Zyme® provides a balanced source of nutrition to support healthy plant growth. Foli-Zyme® assists plants in overcoming the negative impact of stress events. It is suitable for use in a variety of horticulture and broadacre crops

For best results spray when crop is actively growing, after irrigation or natural rainfall. Spray early in the morning or late in the afternoon. Midday sprays may not be effective because of excessive moisture loss.

This product may be applied to the soil through fertigation. Refer to you Corteva Agriscience representative for application rates.

COMPATIBILITY

Prior to tank mixing, always conduct a “Jar Test” using products in proper proportion in order to establish physical compatibility.

APPLICATION

Ensure even coverage of the plant.

STORAGE AND DISPOSAL

Avoid release to environment.

We recommend storing drums a maximum of three drums high. Do not stack pallets. Store container tightly closed in a cool well-ventilated dry place protected from direct sunlight. Do not contaminate water, food or feed by storage or disposal.

™® Trademarks of Corteva Agriscience and its affiliated companies. © 2026 Corteva

20 L

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. Do not dispose of undiluted product on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately.

1,000 L

Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted product on site. Close all valves and arrange for collection under the relevant return program.

SAFETY WARNING

Do not swallow. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin. Users are encouraged to seek further advice.

GHS CLASSIFICATION



- Causes severe skin burns and eye damage.
- May damage fertility or the unborn child.
- Very toxic to aquatic life with long lasting effects.

Obtain special instructions before use.
Do not handle until all safety precautions have been read and understood.
Wash skin thoroughly after handling.
Avoid release to the environment.
Wear protective gloves / protective clothing / eye protection / face protection / hearing protection.

FIRST AID

IF SWALLOWED: Rinse mouth. Do not induce vomiting.
IF ON SKIN (or hair): Take off immediately all contaminated clothing. Rinse skin with water.
IF INHALED: Remove person to fresh air and keep comfortable for breathing. Immediately call a POISON CENTRE / doctor.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTRE / doctor.
IF exposed or concerned: Get medical advice / attention.
Wash contaminated clothing before re-use.
Collect spillage.

Poisons Information Centre Australia 13 11 26.

SAFETY DATA SHEET

Additional information is listed on the Safety Data Sheet which is available from Corteva Agriscience on request. Call Corteva Agriscience Customer Service on number 1800 700 096 or visit www.corteva.com.au.

SPILL AND LEAK MANAGEMENT

Do not touch or walk through spilled material. Dam the area and prevent entry into waterways, and drains. Absorb with material such as sand, soil or sawdust. Collect the spilled product and place in a sealable container. Absorb and collect washings and place in the same sealable

™® Trademarks of Corteva Agriscience and its affiliated companies. © 2026 Corteva

container for disposal. Dam the area of large spills and report them to Corteva Agriscience Emergency Services at 1-800 370 754.

CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product.
2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

™® Trademarks of Corteva Agriscience and its affiliated companies. © 2026 Corteva