DANGER

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KEEP OUT OF REACH OF CHILDREN

Foliar Plus

NUTRITIONAL FOUNDATIONS

(10 – 9 – 3 – 0)

A liquid fertiliser formulated with macro and micronutrients to supplement soil applied fertilisers and enhance plant growth.

ANALYSIS	**(% W/V):**	Boron (B)	0.2 %
Nitrogen (N as total)	10.0 %	Cobalt (Co)	0.07 %
Phosphorus (P as total)	9.0 %	Copper (Cu)	0.15 %
Potassium (K as total)	3.0%	Iron (Fe)	0.1 %
Magnesium (Mg as total)	1.0 %	Manganese (Mn)	0.4 %
Sulphur (S as total)	0.4 %	Molybdenum (Mo)	0.01 %
		Zinc (Zn)	1.5 %
		Maximum biuret	0.82 g/kg N

SHAKE WELL BEFORE USING

Pack sizes: 20, 1000 L

Manufactured by:
Stoller Australia Pty Ltd
1 Creswell Road
Largs Bay, SA, 5016
Australia

Distributed by:
Corteva Agriscience Australia Pty Ltd
Level 9, 67 Albert Avenue
Chatswood, NSW, 2067
Australia

EMERGENCY RESPONSE (ALL HOURS)
RING FROM ANYWHERE IN AUSTRALIA
1800 370 754
(LOCAL CALL FEE ONLY)

IN A TRANSPORT EMERGENCY ONLY
DIAL 000
FOR POLICE OR FIRE BRIGADE

This label complies with the National Code of Practice for Fertilizer Description and Labelling.

Lot Number: See packaging
Manufacturing date: See packaging
Expiration Date: See packaging

™® Trademarks of Corteva Agriscience and its affiliated companies. © 2026 Corteva

DIRECTIONS FOR USE

RESTRAINTS

Excessive use of molybdenum (Mo) can be harmful to stock. Plant levels of Mo can be high for 4 weeks after application. It is advisable to keep stock off treated areas for this period.

Molybdenum application may induce copper deficiency in grazing animals. This is most likely to occur on sandy soils low in copper. Check rate and frequency of molybdenum use with appropriate authorities.

Crop	Rate	Comment
Vegetables	3 - 6 L/ha	To supplement soil applied fertilisers, foliar spray at weekly intervals. Start when crops are a few centimetres high. Continue up to 2 weeks before final harvest.
Trees, Fruits and Nuts	3 - 6 L/ha	Apply 5 - 6 times annually beginning with a late dormancy application for vines and nuts.
Citrus	3 - 7 L/ha	Apply 4 times annually with first application at pre-bloom
Grapevines	3 - 6 L/ha	Can be applied early season (small canopy) through to ripening (veraison) if needed.
Field crops	4 - 8 L/ha	Apply when crops reach 4-leaf stage. Repeat every 14 days until crop matures.
Green House	500 mL /200 L	Spray plants for thorough coverage. Repeat every 14 days until crop matures.
Pasture	5 - 9 L/ha	Maintenance applications in Autumn and Spring when the grass is large enough to get good coverage. For higher production environments, apply 3 times a season beginning at early growth. A minimum dilution rate of 1-part Foliar Plus to 20 parts carrier is recommended for aerial spreading. Can be mixed with most insecticides, fungicides and herbicides.
Turf	Use 2 L in 350 L of water for every 4,000 square metres.	Commence applications in Spring. Repeat every 3 - 4 weeks until grass goes into dormancy.
Soil treatment	5 - 30 L/ha	Foliar Plus can be used as a soil treatment through micro-irrigation (drip).

For further information, consult your Corteva Agriscience representative for advice specific to your situation.

GENERAL INSTRUCTIONS

Apply as a foliar spray in a volume of water sufficient for thorough coverage.

Avoid application in high temperatures or conditions of extreme sunlight.

COMPATIBILITY

Prior to tank mixing, always conduct a “Jar Test” using products in proper proportion in order to establish physical compatibility.

APPLICATION

Foliar Plus may be applied by foliar or soil application methods.

STORAGE AND DISPOSAL

Avoid release to environment.

™® Trademarks of Corteva Agriscience and its affiliated companies. © 2026 Corteva

We recommend storing drums a maximum of three drums high. Do not stack pallets. Store container tightly closed in a cool well-ventilated dry place protected from direct sunlight. Do not contaminate water, food or feed by storage or disposal.
20 L

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. Do not dispose of undiluted product on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately.

1000 L
Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted product on site. Close all valves and arrange for collection under the relevant return program.

SAFETY WARNING

Do not swallow. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin. Users are encouraged to seek further advice.

GHS CLASSIFICATIONS



- Causes skin irritation.
- Causes serious eye irritation.
- May cause cancer.
- May damage fertility or the unborn child.
- Toxic to aquatic life with long lasting effects.

Obtain special instructions before use.
Do not handle until all safety precautions have been read and understood.
Wear protective gloves / protective clothing / eye protection / face protection / hearing protection.
Wash skin thoroughly after handling.
Avoid release to the environment.

FIRST AID

If exposed or concerned: Get medical advice / attention.
IF ON SKIN (or hair): Wash with plenty of water. If skin irritation occurs: Get medical advice / attention.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice / attention.
Take off contaminated clothing and wash it before re-use.
Collect spillage.

Phone a Poisons Information Centre Australia 13 11 26.

SAFETY DATA SHEET

Additional information is listed on the Safety Data Sheet which is available from Corteva Agriscience on request. Call Corteva Customer Service on number 1800 700 096 or visit www.corteva.com.au.

SPILL AND LEAK MANAGEMENT

™® Trademarks of Corteva Agriscience and its affiliated companies. © 2026 Corteva

Do not touch or walk through spilled material. Dam the area and prevent entry into waterways, and drains. Absorb with material such as sand, soil or sawdust. Collect the spilled product and place in a sealable container. Absorb and collect washings and place in the same sealable container for disposal. Dam the area of large spills and report them to Corteva Agriscience Emergency Services at 1-800 370 754.

CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product.
2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.



™® Trademarks of Corteva Agriscience and its affiliated companies. © 2026 Corteva