

blank page

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**Apparent** 

**Alternative**

**HERBICIDE**

ACTIVE CONSTITUENT: 525g/L NONANOIC ACID

For non-selective control of seedling and young broadleaf and grass weeds, for suppression of established weeds and perennial species and for control of Moss and Algae according to the directions for use.  
**IMPORTANT: Read this booklet before use.**



APVMA Approval No: 87680/130979

**AIRR APPARENT PTY LTD**  
15/16 Princes Street, Newport NSW 2106  
P 03 5820 8400 | E [enquiries@apparentag.com.au](mailto:enquiries@apparentag.com.au) | [www.apparentag.com.au](http://www.apparentag.com.au)

**RESTRAINTS:**

DO NOT spray if weeds are wet.

DO NOT apply if rainfall is expected before spray deposits dry on plant surfaces.

DO NOT water treated area for 24 hours to achieve maximum results.

DO NOT apply if heavy rains or storms that are likely to cause runoff from soil are forecast within 3 days of application.

DO NOT irrigate to the point of runoff from soil for at least 3 days after application.

DO NOT allow spray to contact or drift onto plants you do not want killed.

**DIRECTIONS FOR USE**

SITUATION	WEEDS	RATE
Paths, Driveways, around sheds Gardens, Amenity horticulture areas, Protected cropping situations, Around Nursery stock. Spot Spraying in Lawns and Turf.	Seedling and young annual and perennial weeds and grasses including fat hen, potato weed, stinking roger, couch grass, chickweed, lambs tongue, peppergrass, capeweed, common storksbill, clover. More advanced or established weeds may only be suppressed and require repeated application.	70 mL per litre of water
Orchards and vineyards Fallow soil		Mix 7 litres per 100 litres water. Apply 900 - 1400 L spray per ha
Gardens, Paths, Rockeries, Tiled Roofs, Walls, Driveways, Around buildings, Spot Spraying in Lawns and Turf	Algae, Lichen, Liverwort, Moss	70 mL per litre of water

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.  
WITHHOLDING PERIOD (WHP): NOT REQUIRED WHEN USED AS DIRECTED.**

**CRITICAL COMMENTS**

Spray to completely cover weed surface. Reapplication at around 7 days may be necessary if regrowth occurs. This product may damage lawn and turf when used as a spot spray. Perennial weed species may require repeated applications to obtain long term control. To control weeds in protected cropping situations, apply on each side of the crop row taking care not to allow spray to contact any of the crop foliage.
Apply as a band spray 400 mm wide on each side of the orchard or vine crop row. Apply at early vegetative stage of weeds. Best results are obtained at 4 - 8 leaf stage when weed canopy is less than 15 cm in height. Established weeds may require more than one application. Repeat spray after 7 days if required. Spray over all weed foliage to ensure full foliage penetration and coverage. Partial foliage penetration or coverage will result in partial control.
Spray to completely cover plant surface. For drives or heavy infestations, clean off by brushing after 3 days and reapply if necessary.

## GENERAL INSTRUCTIONS

Apparent Alternative Herbicide is a natural contact weedkiller containing pelargonic acid (nonanoic acid). Pelargonic acid occurs naturally in plants including geraniums. It acts on the foliage, causing it to quickly dehydrate and collapse. Results are seen as browning and wilting usually within hours of application. To be effective, Apparent Alternative Herbicide must come into contact with the green parts of the plant. Apparent Alternative Herbicide will control many weed species and is non selective.

### MIXING

Full fill spray tank and commence agitation. Slowly add required amount of Apparent Alternative Herbicide. Top up with required quantity of water. Agitate well before commencing spray procedure. Do not mix more than is needed.

### APPLICATION

Ensure that all weed foliage is totally covered with spray as Apparent Alternative Herbicide is a contact spray only. Partial coverage will give only partial control. Apply by ground equipment only using for example boom or spot sprayers. Apply using a coarse droplet spray. Optimum application volumes are related to weed maturity and density. Contact with the weeds is essential for control, as such apply in sufficient water to ensure complete and thorough coverage of foliage.

DO NOT spray porous surfaces without prior testing. May cause discolouration on porous hard surfaces.

### Resistance Warning:

**PRECAUTION:** For professional use only. Not for use by domestic users.

### Re-Entry Period

Do not allow entry into treated areas until spray has dried. If prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate wetlands or watercourses with this product or used containers. Not compatible with integrated pest management (IPM) programs utilising beneficial arthropods. Minimise spray drift to reduce harmful effects on beneficial arthropods in noncrop areas.

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants, crops, cropping lands or pastures.

### STORAGE AND DISPOSAL:

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility.

If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable

vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

**Refillable Containers:** Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage.

### SAFETY DIRECTIONS

Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with the eyes and skin. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and face shield or goggles. When using the prepared spray wear gauntlet-length chemical resistant gloves and face shield or goggles. Wash hands after use. After each day's use wash gloves, face shield or goggles, and contaminated clothing.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126; New Zealand 0800 764 766. If swallowed, do NOT induce vomiting. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

### CONDITIONS OF SALE:

AIRR APPARENT Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on AIRR APPARENT's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of AIRR APPARENT Pty Ltd has any authority to add to or alter these conditions.